



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

From (5) + (7), $2am^2n^2 - 9bn^4 - 2abn^2 = ab^2 + a^2b \dots \dots \dots (9)$.

Subtracting (9) from (8), $2abn^2 = 2ab(b-a)$. Whence $b-a=n^2$.

Factoring (8), $n^2(2am^2 - 3bn^2) = ab(b-a)$. Whence $2am^2 - 3bn^2 = ab$.

Put $2a=n$ and by easy reduction we have $2m^2 = b(6n+1)$. Since $\frac{1}{2}(6n+1)$ cannot be a square $b/2$ must be a square to make an integral. Then $m^2 = b/2(6n+1)$ is $(n^2 + \frac{1}{2}n)/2 \times (6n+1)$. $n=4$ being the only value that will make both factors a square, $m = \sqrt{[(n^2 + \frac{1}{2}n)/2] \times (6n+1)} = 15$.

$\therefore x=241, y=269, z=149, p=329, q=89, r=191$.

PROBLEMS FOR SOLUTION.

ARITHMETIC.

106. Proposed by ELMER SCHUYLER, High Bridge, N. J.

What is the amount of \$1000 at compound interest for 3 years at 6%, if it be compounded every instant?

107. Proposed by R. V. ALLEN, Hooker Station, Ohio.

A barn, $ABCD$, length $AB=b$ feet, width $AD=a$ feet, standing in an open field, has a horse tethered to a point, P , in the side, AB , distance $AP=c$ feet, with a rope R feet long. Over what area can the horse graze?

*** Solutions of these problems should be sent to B. F. Finkel, not later than March 10.

ALGEBRA.

94. Proposed by J. W. YOUNG, Columbus, Ohio.

Solve: $\left[\frac{x^2 + 14x + 1}{p^4 + 14p^2 + 1} \right]^3 = \frac{x(x-1)^4}{p^2(p^2-1)^4}$.

Burnside and Panton's *Theory of Equations*, page 148, ex. 17.

95. Proposed by SYLVESTER ROBINS, North Branch Depot, N. J.

Substitute *numbers* in place of the letters in the following pattern: $\dots \Delta = \sqrt{81^2 a^2 b^2 c^2} = 81abc \dots b^2 + c^2, a^2 + c^2, a^2 + b^2$; and compute the areas and sides of the whole nest of integral, rational triangles.

*** Solutions of these problems should be sent to J. M. Colaw, not later than March 10.

GEOMETRY.

114. Proposed by WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics and Astronomy, Ohio State University, Athens, Ohio.

If a variable ellipse hyperosculate a fixed ellipse at the extremity of the minor axis, the locus of the foci is a circle whose diameter is equal to the radius of curvature.